


**Journal of Human Sport and Exercise online***J. Hum. Sport Exerc.**Official Journal of the Area of Physical Education and Sport.**Faculty of Education. University of Alicante. Spain***ISSN 1988-5202 / DOI: 10.4100/jhse**

An International Electronic Journal

**Volume 5 Number 2 May 2010****Research Article****A REGRESSION EQUATION FOR THE ESTIMATION OF VO<sub>2</sub>MAX IN NEPALESE MALE ADULTS**Pinaki Chatterjee<sup>1,2</sup> , Alok K. Banerjee<sup>2</sup>, Paulomi Das<sup>3,4</sup>, Parimal Debnath<sup>4</sup>

<sup>1</sup>Department of Physiology, Sudha Rustagi College of Dental Sciences and Research, Faridabad, India. <sup>2</sup>Department of Physical Education, University of Kalyani, Kalyani, West Bengal, India. <sup>3</sup>Department of Physiology, Nepalgunj Medical College, Chisapani, Banke, Nepal. <sup>4</sup>Department of Physical Education, Jadavpur University, Kolkata, India


Received: 16 February 2010; received in revised form: 4 April 2010; accepted: 15 April 2010

**ABSTRACT**

The purpose of the study was to validate the applicability of 20-meter multi stage shuttle run test in Nepalese male College students. 40 Nepalese students were recruited for the study. Direct estimation of cardiorespiratory endurance (VO<sub>2max</sub>) comprised treadmill exercise followed by expired gas analysis by scholander micro-gas analyzer whereas VO<sub>2max</sub> was indirectly predicted by the 20-meter multistage shuttle run test. The difference between the mean (SD) VO<sub>2max</sub> values of direct measurement (VO<sub>2max</sub> = 39.29±1.98 ml/kg/min) and 20-meter multi stage shuttle run test (SPVO<sub>2max</sub> = 39.20±2.27 ml/kg/min) was statistically insignificant (p >0.10). Limits of agreement analysis also suggest that the 20-m multistage shuttle run test can be applied for use with the studied population. The results suggest that the application of the present form of 20-meter multistage shuttle run test be justified in the studied population. For better prediction of VO<sub>2max</sub> a new equation has been developed based on present data.

**Key words:** Beep Test, Aerobic Capacity, Maximum oxygen uptake capacity, Cardiovascular Fitness, Nepalese male.

**Reference Data:** Chatterjee P, Banerjee AK, Das P, Debnath P. A regression equation for the estimation of VO<sub>2max</sub> in nepalese male adults. *J. Hum. Sport Exerc.* 2010; 5(2):127-133.

 **Corresponding author.** Pinaki Chatterjee. Reader and HOD, Department of Physiology, Sudha Rustagi College of Dental Sciences and Research, Faridabad, India.  
Phone: 9339958102/ Fax: 0129-2202958  
E-mail: drpinakichattopadhyay@gmail.com  
© 2010 University of Alicante. Faculty of Education.

**DOI:10.4100/jhse.2010.52.03**

## INTRODUCTION

Direct measurement of maximum oxygen uptake ( $\text{VO}_{2\text{max}}$ ) is recognized as the best single index of aerobic fitness (Astrand & Rodahl, 1986). But the test of the direct measurement of cardiorespiratory endurance ( $\text{VO}_{2\text{max}}$ ) itself is difficult, exhausting and often hazardous to perform regardless the type of ergometer used (Fox, 1973). Since the direct testing procedure is rather complicated on larger populations, several indirect running and walking field tests have been developed. Scientists often calculate  $\text{VO}_{2\text{max}}$  with indirect protocols (Das & Bhattacharya, 1995). It has been stated that equations for predicting  $\text{VO}_{2\text{max}}$  indirectly using field tests are very sensitive to populations tested on. Therefore, before applying any indirect protocol for prediction of  $\text{VO}_{2\text{max}}$ , the validity of the test should be established in a particular population. The 20-m multi stage shuttle run (Leger et al., 1988; Leger & Gadoury., 1989) is often used for measurement of aerobic capacity (Wong et al., 2001; Mota et al., 2002; Guerra et al., 2002; Vicente-Rodriguez et al., 2003; Vicente-Rodriguez et al., 2004). Cooper et al. (2004) studied the repeatability and criterion related validity of the 20-m multistage fitness test as a predictor of maximal oxygen uptake in active young men. Suminski et al. (2004) established the validity of the 20-m MST for measuring aerobic fitness of Hispanic youth of 10 to 12 years of age. Chatterjee et al. (2006) and Chatterjee et al. (2008) studied the validity of 20-m MST in junior Taekwondo players and female university students of India. However, studies on the validity and suitability of this test are not done in any of the Nepalese population until now.

Recent study suggests that genders distinct equations provide more accurate prediction of  $\text{VO}_{2\text{max}}$  from 20-m multistage shuttle run test (Stickland et al., 2003). For this reason only male adults were recruited as subjects and not a male and female-pooled population. The present study was undertaken with an objective to assess the applicability of the 20-m MST to predict  $\text{VO}_{2\text{max}}$  in male College students of Nepal and also to develop a regression equation for use with the particular population.

## METHODS

### *Subjects*

40 male College students from different colleges of Nepal were volunteered for the study. They were selected on the basis of random sampling. The subjects are delimited to all over Nepal, as students from every part of the country come to these colleges for study.

The experimental protocol was fully explained to the participants. They had a light breakfast 2 - 3 hours before the test and refrained from any energetic physical activity for that period. The participants had no history of any major disease and did not follow any physical-conditioning program, except from some recreational sports. Their recreational sports include table tennis, badminton and volleyball. They play either of these games on an average twice a week, and for the duration of half an hour to one hour. Apart from this, most of the students have a practice of cycling for 20 minutes to half an hour daily, as their mode of transport from colleges to residence is bicycle. By analyzing their dietary pattern it reveals that – daily energy intake 2,875-3,000 kcal, of which 15-17% from protein and 10-20% from fat, and the rest from carbohydrate. Considering their life style and habitual activity they may be accepted as true representative of the majority of the Nepalese male College students. The tests were demonstrated to the subjects

before actual administration and they agreed to sign a statement of informed consent. All institutional policies concerning the human subjects in research were followed. The ethical approval was taken from the competent authority.

### *Experimental Design*

Maximum oxygen consumption of each subject was determined by both indirect and direct methods at an interval of 4 days by random sequencing. Indirect one in the half of the subjects followed the direct method whereas indirect one was followed by direct in other half of the subjects to avoid any possibility of bias. Subjects were asked to take complete rest at least for half an hour prior to the exercise, so that pulmonary ventilation and pulse rate might come down to steady state (Chatterjee & Chakravarti, 1986).

#### *Indirect measurement of $VO_{2max}$ using the 20 m MST*

Subjects started running back and forth a 20-metre course and touched the 20 meter (m) line at an initial speed of 8.5 kilometer/hour (km/hr). The speed of the shuttle runs got progressively faster (0.5 km/hr every minute), in accordance with a pace dictated by a sound signal on an audio tape. Several shuttle runs made up each stage, and subjects were instructed to keep pace with the signal for as long as possible. When the subjects could no longer follow the pace, the last stage announced was used to predict  $VO_{2max}$  using the equation of Leger and Gadoury (1989).

#### *Direct measurement of $VO_{2max}$*

The subjects walked on a treadmill to warm up at a speed of 4 km/hr at a 4.5 inclination for duration of 5-min (Slonim et al., 1957). Running at a constant speed of 7 km/hr for a maximum duration of 5 min followed this. The inclination gradient was increased successively from 4.5 until the subject was unable to continue the task. In no case did it exceed 7.5 inclinations. The criteria for maximality was exhaustion and withdrawal from running within the scheduled 5 min period, when the heart rate was about their predicted maximum heart rate and when a further increase of inclination did not bring about any significant rise in oxygen uptake (Chatterjee & Chakravarti, 1986).

### *Gas Analysis*

Low resistance high velocity Collin's Triple "J type" plastic valve was used for the collection of gas by open circuit method (Chatterjee & Chakravarti, 1986). The valve was connected with the Douglas Bag (150-liter) and the expired gas was collected in the second minute of the exhausting final workload, if signs of severe exhaustion supervened. No gas collection was made in the first minute of the workload. The expired gas measured in a wet gasometer (Toshniwal, Germany, CAT No. C G 05.10) and the aliquots of gas samples were analyzed in a Scholander micro gas analysis apparatus following the standard procedure (Consolazio et al., 1963).

### *Validity of the results*

Repeatability was investigated where 22 subjects performed the test twice. The results showed non-significant bias between the two applications of the 20-m MST (mean of the difference  $\pm$  standard deviation of the difference =  $-0.1 \pm 0.9$  ml /kg/min;  $t = -0.7$   $p = 0.60$  with 95% limits of agreement).

For direct measurement of  $VO_{2max}$ , results were corroborated with the previous investigation (Das & Bhattacharya, 1995) in the same laboratory involving an identical population using the same protocol. No significant difference ( $p > 0.1$ ) was observed between the results of the two investigations.

### *Statistical Analysis*

Paired t-test, Pearson's product moment correlation, linear regression statistics and Bland and Altman, (1986) approach for limits of agreement were adopted for statistical analysis of the data. Statistical Package for Social Sciences (SPSS) Microsoft windows Release 11.5 version was used for statistical analysis.

## **RESULTS**

Means and standard deviations of physical characteristics, shuttle predicted  $VO_{2max}$  (SPVO<sub>2max</sub>) by the 20-m multi stage shuttle run test and directly measured  $VO_{2max}$  of the participants are presented in the Table 1.


**Table 1.** Physical parameters, predicted and measured  $VO_{2max}$  of the subjects (N=40)

<b>Parameter</b>	<b>Minimum</b>	<b>Maximum</b>	<b>Mean</b>	<b>Std. Deviation</b>
Age (yr)	20.42	24.92	22.72	1.27
Height (cm)	163.00	167.90	165.90	1.28342
Weight (kg)	52.10	56.50	54.03	1.14
$VO_{2max}$ (ml/kg/min)	36.00	44.20	39.29	1.98
SPVO <sub>2max</sub> (ml/kg/min)	35.60	44.60	39.20	2.27
Shuttle run speed (km/hr)	10.50	12.00	11.10	0.38

No significant variation was observed ( $p > 0.10$ ) between the values of directly measured and predicted  $VO_{2max}$ . The mean difference between  $VO_{2max}$  and SPVO<sub>2max</sub> was 0.09 ml/kg/min with 95% confidence interval  $-0.11$  to  $0.29$  ml/kg/min indicating that 20-m MST predict the maximum oxygen uptake capacity by between  $-0.11$  to  $0.29$  ml/kg/min.

## DISCUSSION

Analysis of data by Bland and Altman method of approach for limits of agreement (Bland & Altman, 1986) between  $SPVO_{2max}$  and  $VO_{2max}$  reveals that limits of agreement are  $-1.13$  to  $1.31$ . These are small enough parameter for 20-m MST to be used confidently in place of direct method (Figure 1). Limits of agreement analysis suggest that application of the present form of 20-m MST should be justified for the studied population.



**Figure 1.** Plotting of difference between  $VO_{2max}$  values against their means (Bland and Altman method of approach).

Highly significant correlation ( $r = 0.97$ ,  $p \leq 0.01$ ) existed between the maximal speed of the 20-m MST and  $VO_{2max}$ . The following equation, derived on the basis of present data will better predict the aerobic fitness in male College students of Nepal.

$$Y = -16.961 + 5.068X,$$

Where,

$$Y = VO_{2max} \text{ (ml/kg/min)}$$

$$X = \text{Maximal shuttle run speed (km/hr)}$$

Using the above new equation the limits of agreement between directly measured  $\text{VO}_{2\text{max}}$  and  $\text{SPVO}_{2\text{max}}$  from the 20-m MST are -1.00 to 0.99. The result suggests that better limits of agreement exist between the two methods when this newly developed equation is used for prediction of  $\text{VO}_{2\text{max}}$  from 20-m MST.

## CONCLUSIONS

Therefore, from the present observations it is concluded that the 20-metre multistage shuttle run test is recommended as a valid method to evaluate aerobic fitness in terms of  $\text{VO}_{2\text{max}}$  within male (age 20.42-24.92 yr.) college students of Nepal. The equation developed on the basis of present data is recommended to be used. This is particularly most useful method for regular monitoring of aerobic fitness in the studied population when a large number of subjects are to be evaluated without the help of a well-equipped laboratory, in less expense and within a short period of time.

## REFERENCES

1. ASTRAND PO, RODAHL K. *Textbook of Work Physiology*. New York: McGraw Hill Book Company; 1986. [[Abstract](#)] [[Back to text](#)]
2. BLAND JM, ALTMAN DG. Statistical method for assessing agreement between two methods of clinical measurements. *Lancet*. 1986; 1(8476):307 - 310. [[Full text](#)] [[Back to text](#)]
3. CHATTERJEE P, BANERJEE AK, MAJUMDAR P, CHATTERJEE P. Validity of the 20-m Multi Stage Shuttle Run Test for the Prediction of  $\text{VO}_{2\text{max}}$  in Junior Taekwondo Players of India. *International Journal of Applied Sports Sciences*. 2006; 18(1):1-7. [[Abstract](#)] [[Back to text](#)]
4. CHATTERJEE P, BANERJEE AK, DAS P, DEBNATH P, CHATTERJEE P. Validity of 20-m multi stage shuttle run test for prediction of maximum oxygen uptake in Indian female university students. *Kathmandu University Medical Journal*. 2008; 6(2):176-180. [[Full text](#)] [[Back to text](#)]
5. CHATTERJEE S, CHAKRAVARTI B. Comparative study of maximum aerobic capacity by three ergometries in untrained college women. *Japanese Journal of Physiology*. 1986; 36(1):151 - 162. [[Full text](#)] [[Back to text](#)]
6. CONSOLAZIO CF, JOHNSON RE, PEKORA LJ. Analysis of Gas Samples. In: Samuel J, Hans field S, editors. *Physiological Measurements of Metabolic Function in Man*. 2<sup>nd</sup> ed. New York: McGraw Hill Book Company; 1963, 507-10. [[Abstract](#)] [[Back to text](#)]
7. COOPER SM, BAKER JS, TONG RJ, et al. The repeatability and criterion related validity of the 20-m multistage fitness test as a predictor of maximal oxygen uptake in active young men. *British Journal of Sports Medicine*. 2005; 39(4):19. [[Full text](#)] [[Back to text](#)]
8. DAS SK, BHATTACHARYA G. A comparison of cardiorespiratory fitness in non-athletes and athletes of eastern India. *Indian Journal of Physiol and Allied Sciences*. 1995; 49:16-23. [[Back to text](#)]
9. FOX EL. A simple, accurate technique for predicting maximal aerobic power. *Journal of Applied Physiology*. 1973; 35(6):914 - 16. [[Abstract](#)] [[Back to text](#)]
10. GUERRA S, RIBEIRO JC, COSTA R, DUARTE J, MOTA J. Relationship between cardiorespiratory fitness, body composition and blood pressure in school children.

- Journal of Sports Medicine & Physical Fitness*. 2002; 42(2):207-13. [[Abstract](#)] [[Back to text](#)]
11. LEGER LA, MERCIER D, GADOURY C, LAMBERT J. The multistage 20 metre shuttle run test for aerobic fitness. *Journal of Sports Sciences*. 1988; 6(2):93-101. [[Abstract](#)] [[Back to text](#)]
  12. LEGER L, GADOURY C. Validity of the 20-m shuttle run test with 1 minute Stages to predict  $\text{VO}_{2\text{max}}$  in Adults. *Canadian Journal Sports Sciences*, 1989; 14(1):21-6. [[Abstract](#)] [[Back to text](#)]
  13. MOTA J, GUERRA S, LEANDRO C, PINTO A, RIBEIRO JC, DUARTE JA. Association of maturation, sex, and body fat in cardiorespiratory fitness. *American Journal of Human Biology*. 2002; 14(6):707-12. [[Abstract](#)] [[Back to text](#)]
  14. SLONIM NB, GILLESPIE DG, HAROLD WH. Peak oxygen uptake of healthy young men as determined by a treadmill method. *Journal of Applied Physiology*. 1957; 10:401 - 404. [[Abstract](#)] [[Back to text](#)]
  15. STICKLAND MK, PETERSEN SR, BOUFFARD M. Prediction of maximal aerobic power from the 20-m multi-stage shuttle run test. *Canadian Journal of Applied Physiology*. 2003; 28 (2):272-82. [[Abstract](#)] [[Back to text](#)]
  16. SUMINSKI RR, RYAN ND, POSTON CS, JACKSON AS. Measuring aerobic fitness of Hispanic youth 10 to 12 years of age. *International Journal of Sports Medicine*. 2004; 25(1):61-7. [[Abstract](#)] [[Back to text](#)]
  17. VICENTE-RODRIGUEZ G, DORADO C, PEREZ-GOMEZ J, GONZALEZ-HENRIQUEZ JJ, CALBET JA. Enhanced bone mass and physical fitness in young female handball players. *Bone*. 2004; 35(5):1208-15. [[Abstract](#)] [[Back to text](#)]
  18. VICENTE-RODRIGUEZ G, JIMENEZ-RAMIREZ J, ARA I, SERRANO-SANCHEZ JA, DORADO C, CALBET JA. Enhanced bone mass and physical fitness in prepubescent footballers. *Bone*. 2003; 33(5):853-9. [[Abstract](#)] [[Back to text](#)]
  19. WONG TW, YU TS, WANG XR, ROBINSON P. Predicted maximal oxygen uptake in normal Hong Kong Chinese schoolchildren and those with respiratory diseases. *Pediatric Pulmonology*. 2001; 31(2):126-32. [[Abstract](#)] [[Back to text](#)]